

Johnson & Johnson Consumer Inc.

-PE Product Labels –

- continued

2789

Other information (continued)

■ contains FD-C yellow no. 5 aluminum lake (barium), as a color additive

■ not between 20-25°C (68-77°F) ■ do not use in carbon or white paper or in ken

polyester 80, pregelatinized starch, titanium dioxide

Drug Facts (continued)

5 aluminum lake (barium), FD-C yellow no. 6 aluminum lake, hydroxypropyl cellulose, hypromellose, magnesium stearate, hydroxypropyl cellulose, polyethylene glycol

Questions or comments? call 1-888-237-7777 (toll-free) or 215-273-8755 (toll-free)

3 00450 58225 6

Distributed by:
JOHNSON & JOHNSON
CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA ©J&J 2011

Warning Facts (continued)

for people with psychological or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the Mx (during if you do not know if your prescription drug contains an Mx). Ask a doctor or pharmacist before taking this product.

■ If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble emptying your bladder
- problems such as occurs with smoking asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product, do not exceed the recommended dose

Stop use and ask a doctor if

- nervousness, dizziness, or sleepiness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- ever gets worse or lasts more than 3 days
- new symptoms occur

The use could be dangerous or a serious condition.

If pregnant or breastfeeding, ask a health professional before use

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ Do not use more than an directed dose over a 24-hour period (warning) ■ Take 2 tablets every 4 hours ■ Do not take more than 10 tablets in 24 hours ■ Ask a doctor

children under 12 years and over

Drug Facts

Active ingredients (in each tablet)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 100 mg
Pseudoephedrine HCl 5 mg
Expectorant
Nasal decongestant

Pain reliever/cough suppressant

Purposes

Uses

- temporarily relieve these symptoms due to the common cold:
- nasal congestion
- headache
- sinus congestion and pressure
- throat irritation
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and ease coughs more productively

Warnings

• Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 4 tablets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg acetaminophen in 24 hours.

• With other drugs containing acetaminophen

• Allergic reactions: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- skin rash
- if skin reaction occurs, stop use and seek medical help right away.

• Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription or nonprescription drug that contains a phenylhydrazine inhibitor (such as chlorzoxazone).

Important: Head all product information before using. Keep this box or important information.

The makers of the SUDAFED® family of products are dedicated to helping you and your family reduce head congestion and sinus pressure.

PREVIOUSLY SUDAFED PE[®] PRESSURE + PAIN + COLD NDC 50580-450-01

SUDAFED^{PE}[®]

HEAD CONGESTION

+ FLU *SEVERE*

Acetaminophen, Dextromethorphan HBr, Guaifenesin,
Phenylephrine HCl, Pain Reliever/Fever Reducer,
Cough Suppressant, Expectorant, Nasal Decongestant

SU PE

- SINUS PRESSURE
- SORE THROAT
- CHEST CONGESTION
- HEADACHE
- COUGH

24 TABLETS

NON-DROWSY

30043256

Drug Facts (continued)

Other information

- do not use if blister unit is torn or broken
- each tablet contains: sodium 3 mg
- store between 20-25°C (68-77°F)

Questions or comments? call 1-888-217-2117 (toll-free) or 1-215-273-8755 (collect)

Inactive ingredients

cantharidin wax, croscarmellose sodium, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, titanium dioxide, triacetin

Does Not Contain
Pseudoephedrine

Open from other side

Drug Facts (continued)

Warnings

- Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (2,500 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product
- Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: rash, skin redness, blisters
- If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks
- with any other drug containing acetaminophen (prescription or nonprescription). If you are unsure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks

Directions

- do not take more than directed (see overdose warning)
- adults and children 12 years and over
- do not take more than 10 tablets in 24 hours
- ask a doctor
- children under 12 years

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away: (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Keep out of reach of children.

Use

- If pregnant or breast-feeding, ask a health professional before use.
- These could be signs of a serious condition:
 - cough comes back or occurs with rash or headache that lasts
 - redness or swelling is present
 - new symptoms occur
 - pain, nasal congestion or cough gets worse or lasts more than 7 days
 - evergets worse or lasts more than 3 days
- Stop use and ask a doctor if:
 - nervousness, dizziness, or sleeplessness occur
 - When using this product do not exceed recommended dose
 - Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin or emphysema
 - cough that occurs with too much phlegm (mucus)
 - persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, diabetes
 - trouble urinating due to an enlarged prostate gland
 - heart disease
 - high blood pressure
 - thyroid disease

Ask a doctor before use if you have

- If you have ever had an allergic reaction to this product or any of its ingredients
- MAOI, ask a doctor or pharmacist before taking this product.

Important: Read all product information before using. Keep this box for important information.

Drug Facts

Active ingredients (in each tablet)

- Acetaminophen 325 mg
- Pain reliever/fever reducer
- Expectorant
- Nasal decongestant

Uses

- temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold
- sinus congestion and pressure
- headache
- minor aches and pains
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways
- temporarily reduces fever

Warnings

- Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (2,500 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product
- Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: rash, skin redness, blisters
- If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks
- with any other drug containing acetaminophen (prescription or nonprescription). If you are unsure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks

Directions

- do not take more than directed (see overdose warning)
- adults and children 12 years and over
- do not take more than 10 tablets in 24 hours
- ask a doctor
- children under 12 years

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away: (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Keep out of reach of children.

Use

- If pregnant or breast-feeding, ask a health professional before use.
- These could be signs of a serious condition:
 - cough comes back or occurs with rash or headache that lasts
 - redness or swelling is present
 - new symptoms occur
 - pain, nasal congestion or cough gets worse or lasts more than 7 days
 - evergets worse or lasts more than 3 days
- Stop use and ask a doctor if:
 - nervousness, dizziness, or sleeplessness occur
 - When using this product do not exceed recommended dose
 - Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin or emphysema
 - cough that occurs with too much phlegm (mucus)
 - persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, diabetes
 - trouble urinating due to an enlarged prostate gland
 - heart disease
 - high blood pressure
 - thyroid disease

Ask a doctor before use if you have

- If you have ever had an allergic reaction to this product or any of its ingredients
- MAOI, ask a doctor or pharmacist before taking this product.

Important: Read all product information before using. Keep this box for important information.



Made in Italy
Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Health Care Division
Fort Washington, PA 19034 USA
© J&J 2018

The makers of the SUDAFED® family of products
are dedicated to helping you and your family reduce
head congestion and sinus pressure.

PREVIOUSLY SUDAFED PE® PRESSURE+PAIN+MUCUS NDC 50580-447-01

SUDAFED^{PE}

HEAD CONGESTION + MUCUS

Acetaminophen, Guaifenesin, Phenylephrine HCl,
Pain Reliever/Fever Reducer, Expectorant, Nasal Decongestant



24 TABLETS

- SINUS PRESSURE
- HEADACHE
- CHEST CONGESTION

NON-DROWSY



Page 4 of 35 PageID #:



4B060 MN C1

[illegible]

Drug Facts (continued)	Drug Facts
<p>Uses</p> <ul style="list-style-type: none"> temporarily relieves these symptoms associated with the common cold or flu: <ul style="list-style-type: none"> headache fever sinus pressure nasal congestion minor body aches and pains reduces swelling of the nasal passages temporarily reduces their breathing through the nose 	<p>Active ingredients (in each caplet)</p> <p>Ibuprofen 200 mg NSAID¹</p> <p>Phenylephrine HCl 10 mg..... nasal decongestant</p> <p>acetaminophen 325 mg..... nonnarcotic anti-inflammatory drug</p> <p>Purposes</p> <p>Pain reliever/fever reducer</p>

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

NON-DROWSY



4B060MNC1

lake, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, modified starch, polyethylene glycol, polyvinyl alcohol, powdered cellulose, pregelatinized starch, sodium starch glycolate, talc, titanium dioxide

Questions or comments?

call 1-888-217-2117 (toll-free) or 215-273-8755 (collect)

- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

SUDAFED^{PE}

NDC 50580-239-01

SUDAFED^{PE}

SINUS CONGESTION

Phenylephrine HCl
Nasal Decongestant

Diphenhydramine HCl,
Phenylephrine HCl
Antihistamine,
Nasal Decongestant

DAYTIME

PE

actual size

• NASAL
CONGESTION

12 TABLETS

10 mg each

NIGHTTIME

PE

actual size

• NASAL
CONGESTION
• RUNNY NOSE

8 TABLETS

TOTAL: 20 TABLETS

SUDAFED^{PE} NIGHTTIME

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- | | |
|--|--|
| adults and children
12 years and over | ■ take 1 tablet every 4 hours
■ do not take more than 6 tablets in 24 hours |
| children under
12 years | ask a doctor |

Other information

- store between 20-25° C (68-77° F)
- do not use if blister unit is torn or broken

Inactive ingredients—carnauba wax, FD&C blue no. 1 aluminum lake, magnesium stearate, microcrystalline cellulose, modified starch, polyethylene glycol, polyvinyl alcohol, powdered cellulose, pregelatinized starch, sodium starch glycolate, talc, titanium dioxide

Questions or comments?

call 1-888-217-2117 (toll-free) or 215-273-8755 (collect)

DO NOT USE IF BLISTER UNIT IS TORN OR BROKEN

Does not contain
Pseudoephedrine

SUDAFED^{PE}
Distributed by: JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA ©J&JCI 2018
Daytime: Active ingredient made in Germany
Nighttime: Made in Italy



30042428

SUDAFED^{PE} PREVIOUSLY SUDAFED^{PE} DAY + NIGHT SINUS CONGESTION NDC 50580-239-01

SINUS CONGESTION

Phenylephrine HCl
Nasal Decongestant

Diphenhydramine HCl, Phenylephrine HCl
Antihistamine, Nasal Decongestant

DAYTIME

PE

actual size

• NASAL
CONGESTION

12 TABLETS 10 mg each

NIGHTTIME

PE

actual size

• NASAL
CONGESTION
• RUNNY NOSE

8 TABLETS | TOTAL: 20 TABLETS

The makers of the SUDAFED^{PE} family of products are dedicated to helping you and your family reduce head congestion and sinus pressure.

<p>Drug Facts (continued)</p> <p>■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma <p>Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers</p> <p>When using this product</p> <ul style="list-style-type: none"> ■ do not exceed recommended dose ■ marked drowsiness may occur ■ avoid alcoholic drinks 	<p>Drug Facts (continued)</p> <p>When using this product do not exceed recommended dose</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ nervousness, or sleeplessness occur ■ symptoms do not improve within 7 days or occur with a fever <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p>Directions</p> <p>adults and children</p> <ul style="list-style-type: none"> ■ take 1 tablet every 4 hours ■ do not take more than 6 tablets in 24 hours <p>children under 12 years</p> <p>ask a doctor</p> <p>Other information</p> <ul style="list-style-type: none"> ■ store between 20-25°C (68-77°F) ■ do not use if blister unit is torn or broken <p>Inactive ingredients</p> <p>canaluba wax, D & C yellow no. 10 aluminum</p>
---	---

Important: Read all product information before using. Keep this box for important information.

Do not take the day and night tablets at the same time. Do not take more than a total of 6 tablets in a 24-hour period. Take only as directed.

<p>SUDAFED PE® DAYTIME</p> <p>Drug Facts</p> <table border="1"> <thead> <tr> <th>Active ingredient (in each tablet)</th> <th>Purpose</th> </tr> </thead> <tbody> <tr> <td>Phenylephrine HCl 10 mg.....</td> <td>Nasal decongestant</td> </tr> </tbody> </table> <p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves sinus congestion and pressure ■ temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies <p>Warnings</p> <p>Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland 	Active ingredient (in each tablet)	Purpose	Phenylephrine HCl 10 mg.....	Nasal decongestant	<p>SUDAFED PE® NIGHTTIME</p> <p>Drug Facts</p> <table border="1"> <thead> <tr> <th>Active ingredients (in each tablet)</th> <th>Purpose</th> </tr> </thead> <tbody> <tr> <td>Diphenhydramine HCl 25 mg.....</td> <td>Antihistamine</td> </tr> <tr> <td>Phenylephrine HCl 10 mg.....</td> <td>Nasal decongestant</td> </tr> </tbody> </table> <p>Uses</p> <ul style="list-style-type: none"> ■ temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: <ul style="list-style-type: none"> ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat ■ nasal congestion ■ temporarily relieves these symptoms due to the common cold: <ul style="list-style-type: none"> ■ runny nose ■ sneezing ■ nasal congestion ■ temporarily relieves sinus congestion and pressure <p>Warnings</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ to make a child sleepy ■ with any other product containing diphenhydramine, even one used on skin 	Active ingredients (in each tablet)	Purpose	Diphenhydramine HCl 25 mg.....	Antihistamine	Phenylephrine HCl 10 mg.....	Nasal decongestant
Active ingredient (in each tablet)	Purpose										
Phenylephrine HCl 10 mg.....	Nasal decongestant										
Active ingredients (in each tablet)	Purpose										
Diphenhydramine HCl 25 mg.....	Antihistamine										
Phenylephrine HCl 10 mg.....	Nasal decongestant										

LIFT HERE

Does Not Contain Pseudoephedrine

SUDAFED^{PE}

Active ingredient made in Germany
Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA ©J&JCI 2018

SUDAFED^{PE}



00450581365

Drug Facts

Active ingredient (in each tablet) Phenylephrine HCl 10 mg..... Nasal decongestant

Purpose Nasal decongestant

Uses

- Temporarily relieves sinus congestion and pressure
- Temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Questions or comments?

call 1-888-217-2177 (toll-free) or 215-273-8755 (collect)

Drug Facts (continued)

Directions

adults and children 12 years and over	ask a doctor
children under 12 years	ask a doctor

■ take 1 tablet every 4 hours
■ do not take more than 6 tablets in 24 hours

Other information

- store between 20-25°C (68-77°F)
- do not use if blister unit is torn or broken

Inactive ingredients carnauba wax, D&C yellow no. 10 aluminum lake, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, modified starch, polyethylene glycol, polyvinyl alcohol, powdered cellulose, pregelatinized starch, sodium starch glycolate, talc, titanium dioxide

30042471/
120482

The makers of the SUDAFED[®] family of products are dedicated to helping you and your family reduce head congestion and sinus pressure.

120482

PREVIOUSLY SUDAFED PE[®] CONGESTION

NDC 50580-437-02



SUDAFED^{PE}

SINUS CONGESTION

Phenylephrine HCl, Nasal Decongestant

36 TABLETS

MAXIMUM STRENGTH

- SINUS PRESSURE
- SINUS CONGESTION

10 mg each

NON-DROWSY

120 483



SUDAFED^{PE}

PREVIOUSLY SUDAFED^{PE} PRESSURE + PAIN

NDC 50580-435-01

120 483

120 483

SINUS PRESSURE + PAIN

Acetaminophen, Phenylephrine HCl

Pain Reliever/Fever Reducer, Nasal Decongestant

MAXIMUM STRENGTH

• SINUS PRESSURE + CONGESTION
• SINUS HEADACHE

actual size

24 TABLETS

NON-DROWSY

120 483

120 483

The makers of the SUDAFED^{PE} family of products are dedicated to helping you and your family reduce head congestion and sinus pressure.

120 483

120 483

Drug Facts

Active ingredients (in each tablet)

Acetaminophen 325 mg..... Pain reliever/fever reducer

Phenylephrine HCl 5 mg..... Nasal decongestant

Uses

- temporarily relieves these symptoms associated with hay fever or other respiratory allergies, and the common cold:
 - sinus congestion and pressure
 - headache
 - throat aches and pains
 - nasal congestion
- temporarily reduces fever

Warnings

Read the important information before using this product. The maximum daily dose of this product is 10 tablets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergic alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- blistering
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or medical conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients.

120 483

120 483

Drug Facts (continued)

Ask a doctor before use if you have:

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product do not exceed recommended dose.

Stop use and ask a doctor if:

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

Pregnant or breastfeeding? Ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away: (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed (see overdose warning)
- adults and children 12 years and over: take 2 tablets every 4 hours
- do not take more than 10 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- do not use if blister unit is torn or broken

Inactive ingredients: carnauba wax, FD&C yellow no. 6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, polyethylene glycol, polysorbate 80, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide.

120 483

120 483

SUDAFED^{PE}

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA ©J&JCI 2018

120 483

120 483

Drug Facts (continued)

Questions or comments?
call 1-888-217-2117 (toll-free) or 215-273-8755 (collect)

120 483

120 483

Does Not Contain Pseudoephedrine

120 483

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Purpose

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
 - sinus congestion and pressure
 - helps clear nasal passages
 - promotes nasal and sinus drainage
 - temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours

with other drugs containing acetaminophen

3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts (continued)

if you have ever had an allergic reaction to this product or any of its ingredients

- Ask a doctor before use if you have
- liver disease
 - heart disease
 - high blood pressure
 - thyroid disease
 - diabetes
 - trouble urinating due to an enlarged prostate gland
 - persistent or chronic cough such as occurs with smoking, asthma or emphysema
 - cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

- Stop use and ask a doctor if
- nervousness, dizziness, or sleeplessness occur
 - pain, nasal congestion or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)

adults and children 12 years and over

- take 2 caplets every 4 hours
- swallow whole; do not crush, chew or dissolve
- do not take more than 10 caplets in 24 hours

children under 12 years

ask a doctor

Other information

- store between 20-25°C (68-77°F)

do not use if blister unit is torn or broken

Drug Facts (continued)

Inactive ingredients anhydrous citric acid, carnauba wax, flavors, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, potassium sorbate, powdered cellulose, pregelatinized starch, sodium benzoate, sodium citrate, sodium starch glycolate, sucralose

Questions or comments? call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

How can we help?
1-877-895-3665





NDC 50580-515-02

TYLENOL[®]

FOR ADULTS

COLD MAX

Acetaminophen,

Dextromethorphan HBr, Phenylephrine HCl
Pain Reliever-Fever Reducer, Cough Suppressant,
Nasal Decongestant



DAY NON-DROWSY

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION



Actual Size

24 CAPLETS

Important: Read all product information before using. Keep this box for important information.

Distributed by:

JOHNSON & JOHNSON CONSUMER INC.


McNeil Consumer Healthcare Division

Fort Washington, PA 19034 USA

©J&JCI 2020



30047051/
123664



Drug Facts

Active ingredients (in each 15 mL) Purpose

Acetaminophen 325 mg Pain reliever/fever reducer
 Dextromethorphan HBr 10 mg.... Cough suppressant
 Phenylephrine HCl 5 mg..... Nasal decongestant

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
 - sinus congestion and pressure
- helps clear nasal passages
- promotes nasal and sinus drainage
- temporarily reduces fever

TYLENOL[®]

COLD + FLU + COUGH

How can we help?

1-877-895-3665

SNAP THIS CODE OR TEXT TYCOLD10 TO 87715

MSG & DATA RATES MAY APPLY
 TEXT HELP FOR HELP
 TEXT STOP FOR STOP



Acetaminophen (325 mg per 15 mL)
 Dextromethorphan HBr (10 mg per 15 mL)
 Phenylephrine HCl (5 mg per 15 mL)

8 fl oz (240 mL)

Made in Canada

Distributed by:

JOHNSON & JOHNSON CONSUMER INC.
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA
 © J&JCI 2018

30042164

Drug Facts (continued)

sorbitol solution, sucralose

Questions or comments?

call 1-877-895-3665 (toll-free) or
 215-273-8755 (collect)

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use

Drug Facts (continued)

and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or

Drug Facts (continued)**Other information**

- each 15 mL contains: sodium 5 mg
- store between 20-25°C (68-77°F). Do not refrigerate.
- do not use if neck band imprinted with "TYLENOL COLD + FLU + COUGH" or foil inner seal imprinted with "SAFETY SEAL®" is broken or missing

Inactive ingredients

alcohol, anhydrous citric acid, FD&C yellow no. 6, flavors, glycerin, propylene glycol, purified water, sodium benzoate,

Drug Facts (continued)**Directions**

- do not take more than directed (see overdose warning)
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

adults and children 12 years and over

- take 30 mL in the dosing cup provided every 4 hours while symptoms last
- do not take more than 150 mL in 24 hours, unless directed by a doctor

children under 12 years

ask a doctor

Drug Facts (continued)

- emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an

Drug Facts (continued)

- enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough

Drug Facts (continued)

- gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

30040371

NDC 50580-369-01

TYLENOL[®]

FOR ADULTS

COLD + FLU + COUGH

Acetaminophen, Dextromethorphan HBr, Doxylamine Succinate, Phenylephrine HCl,
Pain Reliever—Fever Reducer, Cough Suppressant, Antihistamine, Nasal Decongestant



- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- RUNNY NOSE



NIGHT



WILD BERRY
BURST



Alcohol 0.7%

8 fl oz (240 mL)

OPEN TO READ DRUG FACTS
(Warnings, Directions...)




PEEL



30040372

Do not use if neck band imprinted with
"TYLENOL COLD+FLU+ COUGH" or foil
inner seal imprinted with "SAFETY
SEAL[®]" is broken or missing.

**DO NOT USE WITH OTHER MEDICINES
CONTAINING ACETAMINOPHEN**



3 0045-0562-081

Drug Facts

Active ingredients Purpose (in each 15 mL)

Acetaminophen 325 mg Pain reliever/
fever reducer

Dextromethorphan HBr 10 mg... Cough suppressant

Doxylamine Succinate 6.25 mg.....Antihistamine

Phenylephrine HCl 5 mg.....Nasal decongestant

Uses

■ temporarily relieves these common cold/flu symptoms:

- minor aches and pains
- headache
- nasal congestion
- runny nose and sneezing
- cough
- sinus congestion and pressure
- helps clear nasal passages
- sore throat

Drug Facts (continued)

- relieves cough to help you sleep
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional

Drug Facts (continued)

conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as

Drug Facts (continued)

Drug Facts (continued)

- emphysema or chronic bronchitis
- glaucoma
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dose**
- excitability may occur, especially in children
- marked drowsiness may occur

- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
 - pain, nasal congestion or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

Drug Facts (continued)

adults and children 12 years and over	<ul style="list-style-type: none"> ■ take 30 mL in the dosing cup provided every 4 hours while symptoms last ■ do not take more than 150 mL in 24 hours, unless directed by a doctor
children under 12 years	ask a doctor

Other information

- each 15 mL contains: **sodium 5 mg**
- store between 20-25°C (68-77°F). Do not refrigerate.
- **do not use if neck band imprinted with "TYLENOL COLD+FLU+COUGH" or foil inner seal imprinted with "SAFETY SEAL®"**

Drug Facts (continued)

is broken or missing

Inactive ingredients

acetic acid, alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no.

TYLENOL®
COLD + FLU + COUGH
NIGHT

How can we help?
1-877-895-3665

40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Questions or comments?

call **1-877-895-3665** (toll-free) or
215-273-8755 (collect)

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

**SNAP THIS CODE
OR TEXT TYCOLD7
TO 87715**

MSG & DATA RATES MAY APPLY
TEXT **HELP** FOR HELP.
TEXT **STOP** FOR STOP.



Acetaminophen
(325 mg per 15 mL)
Dextromethorphan HBr
(10 mg per 15 mL)
Doxylamine succinate
(6.25 mg per 15 mL)
Phenylephrine HCl
(5 mg per 15 mL)
8 fl oz (240 mL)

Distributed by:


JOHNSON & JOHNSON CONSUMER INC.

McNeil Consumer Healthcare Division

Fort Washington, PA 19034 USA

© J&JCI 2017

30040372



Drug Facts

Active ingredients

Purpose

Drug Facts (continued)

■ temporarily reduces fever

Active ingredients Purpose (in each 30 mL)

Acetaminophen 650 mg.....Pain reliever/
fever reducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Guaifenesin 400 mg.....Expectorant
Phenylephrine HCl 10 mg.....Nasal decongestant

Uses

- temporarily relieves the following cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression,

Drug Facts (continued)

psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

Stop use and ask a doctor if

Drug Facts (continued)

- redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

away. (1-800-222-1222)
Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not take more than directed (see overdose warning)
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

adults and children 12 years and over	<ul style="list-style-type: none"> ■ take 30 mL in the dosing cup provided every 4 hours while symptoms last ■ do not take more than 150 mL in 24 hours, unless directed by a doctor
children under 12 years	ask a doctor

Drug Facts (continued)

Other information

- each 30 mL contains: sodium 10 mg
- store between 20-25°C (68-77°F).
- do not use if neck band imprinted with "WARMING SENSATION" or foil inner seal imprinted with "TYLENOL" is broken or missing

Inactive ingredients

anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, FD&C yellow no. 6, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Drug Facts (continued)

Questions or comments?

call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

TYLENOL®
COLD + FLU SEVERE


How can we help?
1-877-895-3665

Acetaminophen
(650 mg per 30 mL)
Dextromethorphan HBr
(20 mg per 30 mL)
Guaifenesin
(400 mg per 30 mL)
Phenylephrine HCl
(10 mg per 30 mL)

8 fl oz (240 mL)

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
© J&JCI 2021

30049845





JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
© J&JCI 2021 30049845

Drug Facts

Active ingredients Purpose (in each 30 mL)

Acetaminophen 650 mg.....Pain reliever/
fever reducer
Dextromethorphan HBr 20 mg.....Cough suppressant
Guafenesin 400 mg.....Expectorant
Phenylephrine HCl 10 mg.....Nasal decongestant

Uses

- temporarily relieves the following cold/flu symptoms:
 - minor aches and pains
 - headache
 - nasal congestion
 - sore throat
 - cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

Drug Facts (continued)

- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks every day while using this product
- Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
 - blisters
 - rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression,

Drug Facts (continued)

psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

Drug Facts (continued)

- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not take more than directed (see overdose warning)
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

adults and children 12 years and over	<ul style="list-style-type: none"> take 30 mL in the dosing cup provided every 4 hours while symptoms last do not take more than 150 mL in 24 hours, unless directed by a doctor
children under 12 years	ask a doctor

Drug Facts (continued)

Other information

- each 30 mL contains: sodium 10 mg
- store between 20-25°C (68-77°F).
- do not use if neck band imprinted with "WARNING SENSATION" or foil inner seal imprinted with "TYLENOL" is broken or missing

Inactive ingredients

anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, FD&C yellow no. 6, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Drug Facts (continued)

Questions or comments?

call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

OPEN TO READ DRUG FACTS → **PEEL**
(Warnings, Directions...)

Do not use if neck band imprinted with "TYLENOL COLD MAX" or foil inner seal imprinted with "SAFETY SEAL®" is broken or missing.

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN



3 0045-0269-08 9

TYLENOL®
COLD MAX

How can we help?
1-877-895-3665

SNAP THIS CODE
OR TEXT TYCOLD7
TO 87715

MSG & DATA RATES MAY APPLY
TEXT HELP FOR HELP.
TEXT STOP FOR STOP.



Made in Canada
Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
© J&JCI 2018

Acetaminophen (325 mg per 15 mL)
Dextromethorphan HBr (10 mg per 15 mL)
Doxylamine succinate (6.25 mg per 15 mL)
Phenylephrine HCl (5 mg per 15 mL)

8 fl oz (240 mL)

30042166

Drug Facts

Active ingredients (in each 15 mL) Purpose

Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough suppressant
Doxylamine succinate 6.25 mg Antihistamine
Phenylephrine HCl 5 mg Nasal decongestant

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - runny nose and sneezing
 - cough
 - sinus congestion and pressure

Drug Facts (continued)

- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take
- more than 4,000 mg of acetaminophen in 24 hours
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks every day while using this product
- Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening

Drug Facts (continued)

- blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
- Do not use**
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 - if you are now taking a prescription

Drug Facts (continued)

monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

Drug Facts (continued)

- prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

Drug Facts (continued)

- especially in children
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged

2812

- taking sedatives or tranquilizers
- When using this product**
- **do not exceed recommended dose**
 - excitability may occur,

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

Drug Facts (continued)

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed** (see overdose warning)
- mL = milliliter
- use only enclosed dosing cup

Drug Facts (continued)

designed for use with this product. Do not use any other dosing device.

adults and children 12 years and over	<ul style="list-style-type: none"> ■ take 30 mL in the dosing cup provided every 4 hours while symptoms last ■ do not take more than 150 mL in 24 hours, unless directed by a doctor
children under 12 years	ask a doctor

Other information

- each 15 mL contains: sodium 5 mg
- store between 20-25°C (68-77°F). Do not refrigerate.
- **do not use if neck band imprinted with "TYLENOL COLD MAX" or**

Drug Facts (continued)

foil inner seal imprinted with "SAFETY SEAL®" is broken or missing

Inactive ingredients

alcohol, anhydrous citric acid, FD&C blue no. 1, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose


Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



123665

NDC 50580-414-04

TYLENOL® COLD + FLU
FOR ADULTS
SEVERE

COLD + FLU SEVERE

DO NOT USE IF BLISTER UNIT IS TORN OR BROKEN

TYLENOL®
FOR ADULTS

**COLD + FLU
SEVERE**

Acetaminophen,

**Dextromethorphan HBr,
Phenylephrine HCl, Guaifenesin
Pain Reliever-Fever Reducer, Cough Suppressant,
Nasal Decongestant, Expectorant**

Acetaminophen,

**Chlorpheniramine Maleate,
Dextromethorphan HBr, Phenylephrine HCl
Pain Reliever—Fever Reducer, Antihistamine,
Cough Suppressant, Nasal Decongestant**



DAY

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- MUCUS + CHEST CONGESTION



**Actual
Size**

16 CAPLETS



NIGHT

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- RUNNY NOSE



Actual
Size

8 CAPLETS

TOTAL 24 CAPLETS



PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

30047050/
123665

How can we help?
1-877-895-3665



Questions or comments? call 1-877-

re, persists for more than 2 days, is accompanied by vomiting, consult a doctor promptly.

open (prescription or nonprescription). If you
 planning, ask a doctor or pharmacist.
 omeprazole inhibitor (MAOI) (certain drugs
 conditions, or Parkinson's disease), or for
 You do not know if your prescription drug
 n acid or taking this product.
 to this product or any of its ingredients

■ high blood pressure **■** thyroid disease
■ enlarged prostate gland
urinary smoking, asthma or empty sexual
fantasy
■ aortic valve prolapse **■** glaucoma
if you are
■ taking sedatives or tranquilizers

- iron
- increase drowsiness
- ☒ operating machinery

■ redness or swelling is present
back or occurs with rash or headache that lasts
3 or lasts more than 7 days
in professional before use.

medical help or contact a Poison Control
ed attention is critical for adults as well
gns or symptoms.

capslets every 4 hours
which do not crush, chew or dissolve
in more than 10 capsules in 24 hours

~~cid, carnauha wax, F&C blue no. 1 aluminum~~

in many cases, such as cornstarch, modified starch, sorbitol, powdered cellulose, pregelatinized starch and starch hydrolyte, sucralose, then inulin dextride

7-895-3665 (toll-free) or 215-273-8755 (co

Drug Facts (continued)

See it, feel warmth: If sore throat is severe, persists for more than 3 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

■ **With any other drug containing acetaminophen** (prescription or over-the-counter), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ **If you are now taking a prescription non-narcotic antidepressant (MMAO)** certain drugs to treat depression, psychiatric or emotional conditions, or Parkinson's disease, or for 2 weeks after stopping the MMAO drug, if you do not know if your prescription drug contains an MMAO, ask a doctor or pharmacist before taking the product.

Information on this product is not intended to replace the advice of a doctor. All trademarks are the property of their respective owners.

Ask a doctor: **What are you?** You have

- liver disease
- diabetes
- heart disease
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if

- nervousness, dizziness, or sleepiness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache at least

These could be signs of a serious condition.

Drug Facts (continued)

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see overdose warning)

12 years and over	<input type="checkbox"/> swallow whole, do not crush, chew or dissolve <input type="checkbox"/> do not take more than 10 caplets in 24 hours
children under 12 years	ask a doctor

Other information

- each capsule contains sodium 3 mg
- store between 20-25°C (68-77°F)
- do not use if blister unit is torn or broken

Questions or comments? call 817-935-5665 toll-free at 1-855-935-5665 or email info@pharmaceuticalintelligence.com

Intend® Cold + Flu Severe Day Made in Italy
 Distributed by: JOHNSON & JOHNSON CONSUMER INC.

Port Washington, PA 19034 USA

©J&JC| 2020



3 ■■■ 00450 55024 ■■■ 8

Important: Read all product information before using. Keep this box for important information.
Do not take the day and night caplets at the same time. Do not take more than a total of 10 caplets in a 24-hour period. Take only as directed.

TYLENOL® COLD + FLU SEVERE NIGHT

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain relief/fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

Uses ■ for the temporary relief of the following cold/flu symptoms:

■ minor aches and pains	■ headache	■ sore throat	■ nasal congestion
■ cough	■ sinus congestion and pressure	■ sneezing and runny nose	
■ helps clear nasal passages		■ relieves cough to help you sleep	
■ temporarily reduces fever			

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours.

■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

TYLENOL® COLD + FLU SEVERE DAY

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

Uses ■ for the temporary relief of the following cold/flu symptoms:

- minor aches and pains ■ headache ■ sore throat
- nasal congestion ■ cough

■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive ■ temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take **more than 4,000 mg of acetaminophen in 24 hours**.

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

LIFT HERE



Drug Facts

Active ingredients (in each caplet)

Acetaminophen 325 mg...	Pain reliever/fever reducer
Guaifenesin 200 mg...	Expectorant
Phenylephrine HCl 5 mg...	Nasal decongestant

Purpose

Uses

- for the temporary relief of the following cold symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks every day while using this product
- Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
 - blisters
 - rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts (continued)

Inactive ingredients

carnauba wax, croscarmellose sodium, flavor, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, sucralose, titanium dioxide, triacetin

Questions or comments? call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Drug Facts (continued)

- If you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

- Stop use and ask a doctor if**
- nervousness, dizziness, or sleeplessness occur
 - pain, nasal congestion or cough gets worse or lasts more than 7 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none"> ■ take 2 caplets every 4 hours ■ swallow whole—do not crush, chew or dissolve ■ do not take more than 10 caplets in 24 hours
children under 12 years	ask a doctor

Other information

- each caplet contains: sodium 3 mg
- store between 20-25°C (68-77°F)
- do not use if blister unit is torn or broken

How can we help?
1-877-895-3665






OPEN TO READ DRUG FACTS
(Warnings, Directions...)

PEEL



30042165 Do not use if neck band imprinted with
"TYLENOL COLD + MUCUS SEVERE" or
foil inner seal imprinted with "SAFETY
SEAL®" is broken or missing.

DO NOT USE WITH OTHER MEDICINES
CONTAINING ACETAMINOPHEN



3 0045-0521-088

TYLENOL®

COLD + MUCUS SEVERE

Drug Facts

Active ingredients (in each 15 mL) Purpose

Acetaminophen 325 mg Pain reliever/
fever reducer

Dextromethorphan HBr 10 mg Cough suppressant

Guaifenesin 200 mg Expectorant

Phenylephrine HCl 5 mg Nasal decongestant

Uses

■ for the temporary relief of the following cold/flu symptoms:

- minor aches and pains
- headache
- nasal congestion
- sore throat
- cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

How can we help?

1-877-895-3665

**SNAP THIS CODE
OR TEXT TYCOLD8
TO 87715**

MSG & DATA RATES MAY APPLY
TEXT HELP FOR HELP.
TEXT STOP FOR STOP.



Acetaminophen
(325 mg per 15 mL)
Dextromethorphan HBr
(10 mg per 15 mL)
Phenylephrine HCl
(5 mg per 15 mL)
Guaifenesin
(200 mg per 15 mL)

8 fl oz (240 mL)

Made in Canada

Distributed by:

JOHNSON & JOHNSON CONSUMER INC.

McNeil Consumer Healthcare Division

Fort Washington, PA 19034 USA

© J&JCI 2018

30042165

Drug Facts (continued)

- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for

Drug Facts (continued)

depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes

Drug Facts (continued)

- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dose

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough

Drug Facts (continued)

gets worse or lasts more than 7 days

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a

Drug Facts (continued)

Directions

- **do not take more than directed (see overdose warning)**

- mL = milliliter

- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

adults
and
children
12 years
and over

- take 30 mL in the dosing cup provided every 4 hours while symptoms last
- do not take more than 150 mL in 24 hours, unless

Poison Control Center right away.
(1-800-222-1222) Quick medical
attention is critical for adults as well as
for children even if you do not notice
any signs or symptoms. ▶

	directed by a doctor
children under 12 years	ask a doctor

▶

Drug Facts (continued)

Other information

- each 15 mL contains: sodium 5 mg
- store between 20-25°C (68-77°F).
Do not refrigerate.
- do not use if neck band
imprinted with "TYLENOL COLD
+ MUCUS SEVERE" or foil inner
seal imprinted with "SAFETY
SEAL®" is broken or missing

Inactive ingredients

alcohol, anhydrous citric acid,
FD&C blue no. 1, flavor, glycerin,
propylene glycol, purified water,
sodium benzoate, sorbitol solution,
sucralose ▶

Drug Facts (continued)

Questions or comments?

call 1-877-895-3665 (toll-free) or
215-273-8755 (collect)

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



